Access DB# 146/6/

SEARCH REQUEST FORM

Scientific and Technical Information Center

SCIENTIFIC REFERENCE BR

FEB 2 5 RECD

Pat. & T.M. Office

*********	******	*********
STAFF USE ONLY	Type of Search	Vendors and cost where applicable
Searcher: Ed	NA_Sequence (#)	
Searcher Phone #:	AA Sequence (#)	Dialog
Searcher Location:	Structure (#)	Questel/Orbit
Date Searcher Picked Up:	Bibliographic	Dr.Link
Date Completed: 3-4-05	Litigation	Lexis/Nexis
Searcher Prep & Review Time:	Fulltext	Sequence Systems
Clerical Prep Time:	Patent Family	WWW/Internet
Online Time:	Other	Other (specify)
PTO-1590 (8-01)		

10

Attorney Docket 3z16.57-US-02

Y and Y' comprise, each independently, a (disubstituted)methylene group, such as a (di-aromatic)methylene group, for example, 10H-anthracen-9-ylidene group, 9-fluorenylidenyl group, and diarylmethylene group (e.g. diphenylmethylene group); and

Z is a linking group, such as -(CH₂)_m- where m is an integer between 1 and 20, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR₆ group, a CR₇, or a CR₈R₉ group where R₆, R₇, R₈, and R₉ are, independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.

Non-limiting examples of such charge transport materials have the following formulas:

15

10

15

Attorney Docket 3216.57-US-02

These photoreceptors may be used successfully with both dry toners and liquid toners to produce high quality images. The high quality of the images can be maintained after repeated cycling.


Synthesis Of Charge Transport Materials

The synthesis of the charge transport materials of this invention can be prepared by the following multi-step synthetic procedures, although other suitable procedures can be used by a person of ordinary skill in the art based on the disclosure herein.

The first step is the reaction of a (disubstituted)ketone, such as diphenyl ketone, 9-fluorenone, and 10H-anthracenone, with an excess of hydrazine to form the corresponding (disubstituted)ketone hydrazone. In the second step, the (disubstituted)ketone hydrazone reacts with an aromatic aldehyde having a (disubstituted)amine group and a hydroxyl group to form the corresponding azine compound having a (disubstituted)amine group and a hydroxyl group. This step supplies the Y, X, R₁ and R₂ groups or the Y', X', R₃ and R₄ groups of formula (1).

Attorney Docket 3216.57-US-02

- 1 19. The electrophotographic imaging process of claim 18 wherein said organophotoreceptor further comprises a second charge transport material.
 - 20. The electrophotographic imaging process of claim 18 wherein Z has the formula -(CH₂)_m- where m is an integer between 1 and 20, inclusive, and one or more of the methylene groups is optionally replaced by O, S, N, C, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, urethane, urea, an ester group, an NR₆ group, a CR₇, or a CR₈R₉ group where R₆, R₇, R₈, and R₉ are, independently, a bond, H, hydroxyl, thiol, carboxyl, an amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group, or part of a ring group.
 - 21. The electrophotographic imaging process of claim 18 wherein said toner comprises colorant particles.
 - 22. A charge transport material having the formula



where R₁, R₂, R₃, and R₄ comprise, each independently, an alkyl group, an alkenyl group, an aromatic group, a heterocyclic group, or a part of a ring group;

X and X' comprise, each independently, an aromatic group;

Y and Y' comprise, each independently, a (disubstituted)methylene group; and

7 Z is a linking group.

23. The charge transport material of claim 22 wherein X and X' are, each independently, a C₆H₃ group.



Attorney Docket 3216.57-US-02

(24.) The	charge	transport	material	of	claim	22	wherein	the
(disubstitute	d)methyle	ne group is	selected fo	rm th	ne group	consi	sting of a	10H-
anthracen-9-	ylidene gr	oup, a 9-fluo	orenylidene	group	o, and a	diaryln	nethylene	group

25. The charge transport material of claim 22 wherein Z has the formula -(CH ₂) _m -
where m is an integer between 1 and 20, inclusive, and one or more of the methylene
groups is optionally replaced by O, S, N, C, B, P, C=O, O=S=O, a heterocyclic group, an
aromatic group, urethane, urea, an ester group, an NR6 group, a CR7, or a CR8R9 group
where R ₆ , R ₇ , R ₈ , and R ₉ are, independently, a bond, H, hydroxyl, thiol, carboxyl, an
amino group, an alkyl group, an alkenyl group, a heterocyclic group, an aromatic group,
or part of a ring group.

=> file reg FILE 'REGISTRY' ENTERED AT 11:37:49 ON 04 MAR 2005 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT. PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2005 American Chemical Society (ACS)

=> d his

FILE 'LREGISTRY' ENTERED AT 11:15:52 ON 04 MAR 2005 L1 STR

FILE 'REGISTRY' ENTERED AT 11:28:35 ON 04 MAR 2005

L2 0 S L1

L3 STR L1

L4 9 S L3

L5 1778 S L3 FUL

SAV L5 DOT039/A

L6 1 S L1 SSS SAM SUB=L5

L7 19 S L1 SSS FUL SUB=L5 SAV L7 DOT039A/A

FILE 'CAOLD' ENTERED AT 11:35:08 ON 04 MAR 2005 L8 0 S L7

FILE 'ZCAPLUS' ENTERED AT 11:37:38 ON 04 MAR 2005 L9 6 S L7

FILE 'REGISTRY' ENTERED AT 11:37:49 ON 04 MAR 2005

=> d 17 que stat

NODE ATTRIBUTES:


NSPEC IS RC ATIS RC ΑT 11 **NSPEC** IS RC AT 18 **NSPEC NSPEC** IS RC AT 19 CONNECT IS M3 RC AT 4 CONNECT IS M3 RC AT DEFAULT MLEVEL IS ATOM 4 **GGCAT** IS UNS AT12 **GGCAT** IS UNS ΑT DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED NUMBER OF NODES IS 12

STEREO ATTRIBUTES: NONE

L3 STR



NODE ATTRIBUTES:

5 AΤ NSPEC IS RC NSPEC IS RC AT 11 NSPEC IS RC AT 18 ΑT 19 **NSPEC** IS RC CONNECT IS M3 RC AT 4 . RC AT 12 CONNECT IS M3 DEFAULT MLEVEL IS ATOM GGCAT IS UNS AT **GGCAT** IS UNS AT 12 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED NUMBER OF NODES IS 12

STÈREO ATTRIBUTES: NONÈ

L5 1778 SEA FILE=REGISTRY SSS FUL L3

L7 19 SEA FILE=REGISTRY SUB=L5 SSS FUL L1

100.0% PROCESSED 1729 ITERATIONS

19 ANSWERS

SEARCH TIME: 00.00.01

=> file zcaplus FILE 'ZCAPLUS' ENTERED AT 11:40:20 ON 04 MAR 2005 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT. PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

=> d 19 1-6 all hitstr

L9 ANSWER 1 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN

AN 2005:1965 ZCAPLUS

DN 142:103066

ED Entered STN: 31 Dec 2004

TI Azine-based dimeric charge transport materials

IN Tokarski, Zbigniew; Jubran, Nusrallah; Getautis, Vytautas; Gaidelis, Valentas; Daskeviciene, Maryte; Montrimas, Edmundas; Paulauskaite, Ingrida; Sidaravicius, Jonas

PA USA

SO U.S. Pat. Appl. Publ., 20 pp.

CODEN: USXXCO

DT Patent

LA English

IC ICM G03G005-06 ICS C07C251-72

NCL 430058350; 430072000; 430077000; 430074000; 430058650; 564251000

CC 74-3 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

FAN.CNT 1

11111.	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
ΡΙ	US 2004265717	A1	20041230	US 2004-760039	200401
	EP 1494080	A1	20050105	EP 2004-253868	16 200406 29

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC,

PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR 20050127 JP 2004-194403 JP 2005025192 A2 200406 30 Ρ 20030630 PRAI US 2003-483726P US 2004-760039 Α 20040116 CLASS CLASS PATENT FAMILY CLASSIFICATION CODES PATENT NO. US 2004265717 ICM G03G005-06 C07C251-72 ICS 430058350; 430072000; 430077000; 430074000; NCL 430058650; 564251000 C07C251/88 US 2004265717 ECLA EP 1494080 C07C251/88 ECLA JP 2005025192 FTERM 2H068/AA20; 2H068/AA34; 2H068/AA35; 2H068/AA54; 2H068/AA55; 2H068/BA16; 2H068/BA18; 2H068/BA23; 4C036/AD08; 4C036/AD20; 4H006/AA01; 4H006/AA03; 4H006/AB76; 4H006/TA04; 4H006/TB14; 4H006/TB36; 4H006/TB76

$$Y = N - N =$$
 $R^{1} - N - X - Z - X^{1} - N - R^{1}$
 R^{2}
 R^{2}

GΙ

Improved organo photoreceptor comprises an elec. conductive AB substrate and a photoconductive element on the elec. conductive substrate, the photoconductive element comprising: (a) a charge transport material having the formula I (R1-4 = alkyl group, alkenyl group, arom. group, heterocyclic group, or a part of a ring group; X and X' = arom. group; Y and Y' = (disubstituted) methylene group; and Z is a linking group); (b) a charge generating compd.; and (c) an elec. conductive substrate on which said charge transport material and said charge generating compd. are located. Corresponding electrophotog. apparatuses and imaging methods are also described. azine dimeric electrophotog photoreceptor charge transport material ST ΙT Electrophotographic photoconductors (photoreceptors) (azine-based dimeric charge transport materials)

IT 816463-93-1P 816463-94-2P 816463-95-3P 816463-96-4P 816463-97-5P 816463-98-6P 816463-99-7P 816464-00-3P 816464-01-4P

816464-02-5P

(azine-based dimeric charge transport materials for electrophotog.)

IT 2915-84-6P, 2,7-Diamino-9-fluorenone 122010-64-4P 215377-16-5P 816464-03-6P 816464-04-7P 816464-05-8P 816464-07-0P 816464-08-1P

(prepn. of azine-based dimeric charge transport materials for electrophotog.)

1T 80-05-7, reactions 90-93-7 106-89-8, Epichlorohydrin, reactions 108-46-3, 1,3-Benzenediol, reactions 486-25-9, 9-Fluorenone 540-63-6, 1,2-Ethanedithiol 626-04-0, 1,3-Benzenedithiol 1072-71-5, 1,3,4-Thiadiazolidine-2,5-dithione 2425-79-8, 1,4-Butanediol diglycidyl ether 17754-90-4, 4-Diethylaminosalicylaldehyde 19362-77-7, 4,4'-Thiobisbenzenethiol 31551-45-8, 2,7-Dinitro-9-fluorenone

(prepn. of azine-based dimeric charge transport materials for electrophotog.)

IT 13629-22-6P 816464-06-9P

(prepn. of azine-based dimeric charge transport materials for electrophotog.)

IT 816463-93-1P 816463-94-2P 816463-95-3P 816463-96-4P 816463-97-5P 816463-98-6P 816463-99-7P 816464-00-3P 816464-01-4P 816464-02-5P

(azine-based dimeric charge transport materials for electrophotog.)

- RN 816463-93-1 ZCAPLUS
- CN Benzaldehyde, 2,2'-[1,2-ethanediylbis[thio(2-hydroxy-3,1-propanediyl)oxy]]bis[4-(diethylamino)-, bis(9H-fluoren-9-ylidenehydrazone) (9CI) (CA INDEX NAME)

PAGE 1-B

RN 816463-94-2 ZCAPLUS

CN Benzaldehyde, 2,2'-[1,3-phenylenebis[oxy(2-hydroxy-3,1-propanediyl)oxy]]bis[4-(diethylamino)-, bis(9H-fluoren-9-ylidenehydrazone) (9CI) (CA INDEX NAME)

PAGE 1-B

RN 816463-95-3 ZCAPLUS

CN Benzaldehyde, 2,2'-[1,4-butanediylbis[oxy(2-hydroxy-3,1-propanediyl)oxy]]bis[4-(diethylamino)-, bis(9H-fluoren-9-

ylidenehydrazone) (9CI) (CA INDEX NAME)

PAGE 1-A

PAGE 1-B

__NEt2

RN 816463-96-4 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

PAGE 1-B

RN 816463-97-5 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

PAGE 1-B

RN 816463-98-6 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

RN 816463-99-7 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

RN 816464-00-3 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

RN 816464-01-4 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

-NEt2

___NEt2

RN 816464-02-5 ZCAPLUS CN INDEX NAME NOT YET ASSIGNED

L9 ANSWER 2 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN

AN 2004:231790 ZCAPLUS

DN 141:63806

ED Entered STN: 22 Mar 2004

TI Regioselective HON-addition of bifunctional hydrazone oximes to Pt(IV)-bound nitriles

AU Garnovskii, Dmitrii A.; Pombeiro, Armando J. L.; Haukka, Matti; Sobota, Piotr; Kukushkin, Vadim Yu.

CS Centro de Quimica Estrutural, Complexo I, Instituto Superior Tecnico, Lisbon, 1049-001, Port.

SO Dalton Transactions (2004), (7), 1097-1103 CODEN: DTARAF; ISSN: 1477-9226

PB Royal Society of Chemistry

DT Journal

LA English

CC 78-7 (Inorganic Chemicals and Reactions) Section cross-reference(s): 75

OS CASREACT 141:63806

Treatment of trans-[PtCl4(RCN)2] (R = Me, Et) with the hydrazone oximes MeC(:NOH)C(R'):NNH2 (R' = Me, Ph) at 45.degree. in CH2Cl2 gave trans-[PtCl4{NH:C(R)ON:C(Me)C(R'):NNH2}2] (R/R' = Me/Ph 1, Et/Me 2, Et/Ph 3) due to the regioselective OH-addn. of the bifunctional MeC(:NOH)C(R'):NNH2 to the nitrile group. The reaction of 3 and Ph3P:CHCO2Me gave the Pt(II) complex trans-[PtCl2{NH:C(Et)ON:C(Me)C(Ph):NNH2}2] (4). In 4, the imine ligand was liberated by substitution with 2 equiv of dppe (bis(1,2-diphenylphosphino)ethane) in CDCl3 to give, along with the free ligand, [Pt(dppe)2]Cl2. The free iminoacyl hydrazone, having a

restricted life-time, decomps. at 20-25.degree. in .apprx.20 h to the parent organonitrile and the hydrazone oxime. The Schiff condensation of the free NH2 groups of 4 with arom. aldehydes, i.e. 2-OH-5-NO2-benzaldehyde and 4-NO2-benzaldehyde, brings about the formation of the Pt(II) complexes trans- $[PtC12{NH:C(Et)ON:C(Me)C(Ph):NN:CH(C6H3-2-OH-5-NO2)}2]$ (5) and trans-[PtCl2{NH:C(Et)ON:C(Me)C(Ph):NN:CH(C6H4-4-NO2)}2] (6), resp., contg. functionalized remote peripheral groups. Metalization of 5, which can be considered as a novel type of metallo-ligand, was achieved by its reaction with M(OAc) 2.cntdot.nH2O (M = Cu, n = 2; M = Co, n = 4) in a 1 : 1 molar ratio furnishing solid heteronuclear compds. [Pt]:[M] = 1 : 1. The complexes were characterized by C, H, N elemental analyses, FAB+ mass-spectrometry, IR, 1H, 13C{1H} and 195Pt NMR spectroscopies; x-ray structures were detd. for 3, 4 and platinum hydrazoneoxime nitrile prepn; crystal structure platinum hydrazoneoxime nitrile Crystal structure Molecular structure (of platinum complexes with bifunctional ligands derived from hydrazone oximes and nitriles) 107-12-0, Propanenitrile (formation from decomp. of iminoacyl hydrazone) 17116-21-1, Bis[1,2-bis(diphenylphosphino)ethane]platinum(2+) dichloride (formation from reaction of dppe with platinum complexes contg. bifunctional ligands derived from hydrazone oximes and nitriles) 709046-49-1 (formation from reaction of dppe with platinum complexes contg. bifunctional ligands derived from hydrazone oximes and nitriles and decomp.) 709046-40-2P 709046-45-7P 709046-41-3P (prepn. of) 709046-46-8P 709046-48-0P (prepn. of polymeric) 709046-43-5P (prepn., crystal structure and reaction with arom. aldehydes) 709046-42-4P (prepn., crystal structure and reaction with phosphorus ylide) 709046-44-6P (prepn., crystal structure and reaction with transition metal acetates) 143729-50-4, trans-Bis(acetonitrile)tetrachloroplatinum 342028-87-9, trans-Tetrachlorobis (propanenitrile) platinum (reaction with hydrazone oximes) 97-51-8, 5-Nitrosalicylaldehyde 555-16-8, 4-Nitrobenzaldehyde, 1663-45-2, 1,2-Bis (diphenylphosphino) ethane

(reaction with platinum complex contq. bifunctional ligands

ST

ΙT

ΙT

ΙΤ

ΙT

ΙT

ΙT

ΙT

ΙΤ

ΙT

ΙT

ΙΤ

derived from hydrazone oximes and nitriles)

6874-04-0 ΙT

(reaction with platinum nitrile complexes)

ΙT 41939-99-5

> (reaction with platinum nitrile complexes and formation from decomp. of iminoacyl hydrazone)

2605-67-6, Methyl (triphenylphosphoranylidene)acetate ΙT

(reducing agent; reaction with platinum complex contg.

bifunctional ligands derived from hydrazone oximes and nitriles) THERE ARE 93 CITED REFERENCES AVAILABLE FOR THIS RECORD 93 RE.CNT RE

- (1) Adams, H; Inorg Chem Commun 2000, V3, P24 ZCAPLUS
- (2) Allen, F; J Chem Soc, Perkin Trans 2 1987, PS1 ZCAPLUS
- (3) Altomare, A; J Appl Crystallogr 1999, V32, P115 ZCAPLUS
- (4) Anderson, G; Inorg Chim Acta 1983, V76, PL251 ZCAPLUS
- (5) Ang, H; J Chem Soc, Dalton Trans 1993, P847 ZCAPLUS
- (6) Bagrov, F; Russ J Org Chem 2001, V37, P15 ZCAPLUS
- (7) Beck, B; Dalton Trans 2003, P2533 ZCAPLUS
- (8) Belinski, J; J Coord Chem 1988, V19, P159 ZCAPLUS
- (9) Bieda, K; Inorg Chem 1993, V32, P4209 ZCAPLUS
- (10) Bogdanovic, G; Polyhedron 2001, V20, P2231 ZCAPLUS
- (11) Bokach, N; Inorg Chem 2003, V42, P836
- (12) Cariati, F; Inorg Chem 2002, V41, P6597 ZCAPLUS
- (13) Chakrabarti, P; Indian J Chem, Sect A 2000, V39, P571
- (14) Charmier, M; Dalton Trans 2003, P2540
- (15) Chen, W; Eur J Inorg Chem 2002, P2664
- (16) Chen, W; Inorg Chim Acta 2003, V342, P88 ZCAPLUS
- (17) Chin, J; Acc Chem Res 1991, V24, P145 ZCAPLUS
- (18) Constable, E; Metals and Ligand Reactivity. An Introduction to the Organic Chemistry of Metal Complexes 1995, P65
- (19) Cotton, F; Polyhedron 1998, V17, P2781 ZCAPLUS
- (20) da Rocha, Z; Adv Chem Ser 1997, P297 ZCAPLUS
- (21) Dinda, R; J Chem Soc, Dalton Trans 2002, P4434 ZCAPLUS
- (22) Dutta, S; Inorg Chem 2002, V41, P5555 ZCAPLUS
- (23) Eglin, J; Comments Inorg Chem 2002, V23, P23 ZCAPLUS
- (24) Endres, H; Comprehensive Coordination Chemistry 1987, V2, P261
- (25) Fang, C; Organometallics 2001, V20, P2525 ZCAPLUS
- (26) Farrugia, L; J Appl Crystallogr 1999, V32, P837
- (27) Fomina, I; J Organomet Chem 2001, V636, P157 ZCAPLUS
- (28) Foster, M; J Chem Soc 1912, P2235
- (29) Freisinger, E; J Chem Soc, Dalton Trans 2000, P3281 ZCAPLUS
- (30) Fun, H; Acta Crystallogr, Sect C 1999, V55, P896
- (31) Galic, N; J Mol Struct 2001, V559, P187 ZCAPLUS
- (32) Grzybowski, J; Inorg Chem 1993, V32, P5266 ZCAPLUS
 (33) Iskander, M; J Coord Chem 2003, V56, P1075 ZCAPLUS
- (34) Iskander, M; Polyhedron 2001, V20, P2787 ZCAPLUS
- (35) John, R; Polyhedron 2002, V21, P2515 ZCAPLUS
- (36) Kim, I; Inorg Chim Acta 2001, V317, P12 ZCAPLUS

- (37) Kitano, K; J Chem Soc, Dalton Trans 2000, P995 ZCAPLUS
- (38) Kogan, V; Russ J Coord Chem 1998, V24, P177 ZCAPLUS
- (39) Kopylovich, M; Inorg Chem 2002, V41, P4798 ZCAPLUS
- (40) Kopylovich, M; Inorg Chem 2003, V42, P7239 ZCAPLUS
- (41) Kopylovich, M; J Chem Soc, Perkin Trans 1 2001, P1569 ZCAPLUS
- (42) Kukushkin, V; WO 01/98283 A1 ZCAPLUS
- (43) Kukushkin, V; Chem Rev 2002, V102, P1771 ZCAPLUS
- (44) Kukushkin, V; Inorg Chem 1998, V37, P6511 ZCAPLUS
- (45) Kukushkin, V; Inorg Chem 2000, V39, P216 ZCAPLUS
- (46) Kukushkin, V; Russ J Inorg Chem (Engl Transl) 1988, V33, P1085
- (47) Kukushkin, V; Zh Neorg Khim 1988, V33, P1905 ZCAPLUS
- (48) Kukushkin, Y; Inorg Chim Acta 1999, V285, P203 ZCAPLUS
- (49) Li, Q; Dalton Trans 2003, P1551 ZCAPLUS
- (50) Lin, S; J Chem Soc, Dalton Trans 2002, P1595 ZCAPLUS
- (51) Luzyanin, K; J Chem Soc, Dalton Trans 2002, P1882 ZCAPLUS
- (52) Makarycheva-Mikhailova, A; Inorg Chem 2003, V42, P2805 ZCAPLUS
- (53) Makarycheva-Mikhailova, A; Inorg Chem 2003, V42, P301 ZCAPLUS
- (54) Mandlik, P; J Indian Chem Soc 2002, V79, P689 ZCAPLUS
- (55) Mandlik, P; Pol J Chem 2003, V77, P129 ZCAPLUS
- (56) Maresca, L; J Am Chem Soc 1986, V108, P1180 ZCAPLUS
- (57) Michelin, R; Coord Chem Rev 1996, V147, P299 ZCAPLUS
- (58) Michelin, R; Inorg Chim Acta 1994, V222, P327 ZCAPLUS
- (59) Mikuriya, M; Chem Lett 1998, P421 ZCAPLUS
- (60) Murahashi, S; Acc Chem Res 2000, V33, P225 ZCAPLUS
- (61) Murray, B; 2003 ZCAPLUS
- (62) Murray, B; Org React Mech 2003, P1 ZCAPLUS
- (63) Otwinowski, Z; Methods Enzymol 1997, V276, P307 ZCAPLUS
- (64) Pal, S; 2002 ZCAPLUS
- (65) Pal, S; Proc Indian Acad Sci, Chem Sci 2002, V114, P417 ZCAPLUS
- (66) Parkins, A; Platinum Met Rev 1996, V40, P169 ZCAPLUS
- (67) Patole, J; Inorg Chim Acta 2001, V318, P207 ZCAPLUS
- (68) Pavlishchuk, V; Chem Commun 2002, P468 ZCAPLUS
- (69) Pavlishchuk, V; Inorg Chem 1999, V38, P1759 ZCAPLUS
 (70) Pombeiro, A; PT 102618 P 2001 ZCAPLUS
- (71) Pombeiro, A; Comprehensive Coordination Chemistry, 2nd edn V1, P585 ZCAPLUS
- (72) Pombeiro, A; Comprehensive Coordination Chemistry, 2nd edn 2003, V1, P639
- (73) Rana, A; Polyhedron 2003, V22, P3075 ZCAPLUS
- (74) Richardson, D; J Biol Inorg Chem 1999, V4, P266 ZCAPLUS
- (75) Rousselet, G; Tetrahedron Lett 1993, V34, P6395 ZCAPLUS
- (76) Sandbhor, U; J Inorg Biochem 2002, V90, P127 ZCAPLUS
- (77) Sheldrick, G; SHELXL-97, Program for Crystal Structure Refinement
- (78) Sheldrick, G; SHELXS97, Program for Crystal Structure Determination 1997
- (79) Sheldrick, G; SHELXTL v 6.12 2002
- (80) Syamala, A; Inorg Chem 1991, V30, P4699 ZCAPLUS

- (81) Szczena, B; New J Chem 2002, V26, P243
- (82) Ukwueze, A; Int J Chem 2003, V13, P181 ZCAPLUS
- (83) Vicente, J; Inorg Chem 2001, V40, P2051 ZCAPLUS
- (84) Voloshin, Y; Inorg Chim Acta 1999, V284, P180 ZCAPLUS
- (85) Voloshin, Y; Inorg Chim Acta 2000, V299, P104 ZCAPLUS
- (86) Voloshin, Y; Polyhedron 2001, V20, P2721 ZCAPLUS
- (87) Voloshin, Y; Russ Chem Bull 2002, V51, P1015 ZCAPLUS
- (88) Voloshin, Z; Inorg Chem Commun 1998, V1, P328
- (89) Wagner, G; Inorg Chem 2001, V40, P1683 ZCAPLUS
- (90) Wagner, G; Inorg Chem 2001, V40, P264 ZCAPLUS
- (91) Wagner, G; J Am Chem Soc 2000, V122, P3106 ZCAPLUS
- (92) Weng Ng, S; Z Kristallogr 1997, V212, P277
- (93) Yilmaz, I; Heteroat Chem 2003, V14, P617 ZCAPLUS
- IT 709046-44-6P

(prepn., crystal structure and reaction with transition metal acetates)

RN 709046-44-6 ZCAPLUS

CN Platinum, dichlorobis[[C(E)]-2-hydroxy-5-nitrobenzaldehyde (2E)-[(2E)-2-[[(1Z)-1-(imino-.kappa.N)propoxy]imino]-1-phenylpropylidene]hydrazone]-, (SP-4-1)- (9CI) (CA INDEX NAME)

PAGE 1-A

PAGE 1-B

L9 ANSWER 3 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN

AN 1990:45614 ZCAPLUS

DATE

198707 17

```
DN
    112:45614
ED
    Entered STN: 04 Feb 1990
ΤI
    Electrophotographic photoreceptor using azo dye
    Shiino, Yasuko; Umehara, Masashige
IN
    Canon K. K., Japan
PA
    Jpn. Kokai Tokkyo Koho, 19 pp.
SO
    CODEN: JKXXAF
DT
    Patent
LA
    Japanese
    ICM G03G005-06
IC
CC
    74-3 (Radiation Chemistry, Photochemistry, and Photographic and
    Other Reprographic Processes)
FAN.CNT 1
    PATENT NO.
                       KIND
                              DATE APPLICATION NO.
     _____
                  A2 19890124 JP 1987-177029
    JP 01021458
PΙ
PRAI JP 1987-177029
                              19870717
CLASS
```

PATENT NO. CLASS PATENT FAMILY CLASSIFICATION CODES

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

- AB In the title photoreceptor, a photoconductive layer contains an azo dye I [R1, R2 = H, alkyl, aralkyl, aryl; Ar1 = aryl, arom. heterocyclic group; R2 and Ar1 may form a ring; X may condense with the benzone ring to form a polyarom. or heterocyclic ring; Ar2 = arom. hydrocarbon ring, arom. heterocyclic ring; n = 1-4]. Efficiency of carrier generation and/or transportation can be improved with the above photoreceptor. A photoreceptor with II showed Vo -700 V and E1/2 lx-s.
- ST azo dye electrophotog photoreceptor

JP 01021458 ICM G03G005-06

IT Dyes, azo

GI

(electrophotog. photoconductive layer contg.)

IT Electrophotographic photoconductors

(photoconductive layer contg. azo dye for)

IT123576-96-5123576-97-6123576-98-7123576-99-8123577-00-4123577-01-5123577-02-6123577-03-7123577-04-8123577-05-9123577-06-0123577-07-1123577-08-2123577-09-3123598-86-7123598-87-8

124633-35-8

(photoconductive layer contg., for electrophotog. photoreceptor) RN 123576-97-6 ZCAPLUS

CN 2-Naphthalenecarboxaldehyde, 4,4'-[(3,3'-dimethoxy[1,1'-biphenyl]-4,4'-diyl)bis(azo)]bis[3-hydroxy-, bis[[(2-chlorophenyl)methylene]hydrazone] (9CI) (CA INDEX NAME)

RN 123577-03-7 ZCAPLUS

CN 2-Naphthalenecarboxaldehyde, 4,4'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)bis(azo)]bis[3-hydroxy-, 2,2'-bis[[(4-methoxyphenyl)methylene]hydrazone] (9CI) (CA INDEX NAME)

PAGE 2-A
OMe

RN 123577-04-8 ZCAPLUS

CN 2-Naphthalenecarboxaldehyde, 4,4'-[2,6-benzoxazolediylbis(azo)]bis[3-hydroxy-, bis[[(2-chlorophenyl)methylene]hydrazone] (9CI) (CA INDEX NAME)

RN 123598-86-7 ZCAPLUS

CN 2-Naphthalenecarboxaldehyde, 4,4'-[1,2-ethenediylbis(4,1-phenyleneazo)]bis[3-hydroxy-, bis[[(2-fluorophenyl)methylene]hydrazone] (9CI) (CA INDEX NAME)

```
L9 ANSWER 4 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN
```

AN 1988:560521 ZCAPLUS

DN 109:160521

ED Entered STN: 28 Oct 1988

TI Electrophotographic photoreceptor using phthalocyanine dye and bishydrazone compound in the photoconductor layer

IN Horie, Seiji; Makino, Naonori; Sato, Hideo

PA Fuji Photo Film Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 13 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM G03G005-04

ICA G03G005-06

CC 74-3 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 63048552	A2	19880301	JP 1986-191774	198608 18

```
US 4814245
                          Α
                                19890321
                                            US 1987-86449
                                                                    198708
                                                                    18
                          Α
                                19860818
PRAI JP 1986-191774
CLASS
 PATENT NO.
                 CLASS PATENT FAMILY CLASSIFICATION CODES
                 ICM
 JP 63048552
                        G03G005-04
                 ICA
                        G03G005-06
     For diagram(s), see printed CA Issue.
GI
     The charge-generating layer contains a phthalocyanine dye (e.g.,
AΒ
     .epsilon.-type Cu phthalocyanaine or AlCl3-phthalocyanine complex),
     and the charge-transporting layer contains .gtoreq.1 bis-hydrazone
     compds. (I) and (II) (R1, R2 = C1-12 \text{ alkyl}, C7-20 \text{ aralkyl},
    monovalent single or .qtoreq. 2-4-ring condensed arom.. hydrocarbon
     residue; R1 and R2 may form heterocycle; R3 = H, C1-12 alkyl, C7-20
     aralkyl, aryl; R4, R7 = C1-12 alkyl, C7-20 aralkyl, aryl, halo,
     alkoxy, aryloxy; R5, R6, R8 = C1-12 alkyl, C7-20 aralkyl, aryl; R5
     and R6 may bond together to form an N-heterocycle; X = (III); l, n =
     0, 1-6; m = 0, 1; Y = 0, S, Se, imino, CH2; and Z = moiety necessary
     to form benzene and naphthalene ring). This electrophotog.
     photoreceptor provides high sensitivity and stability.
     electrophotog photoreceptor phthalocyanine dye bishydrazone
ST
     Electrophotographic photoconductors
ΙT
        (composite, contg. bishydrazone compds. and phthalocyanine dyes)
ΙT
     147-14-8
                14154-42-8
        (charge-generating layer contg., for electrophotog.
        photoreceptor)
     101158-34-3 101158-35-4
                                 101158-37-6
                                               101158-38-7
                                                              101158-40-1
ΙΤ
     101158-41-2
                  101158-43-4
                                 101158-47-8
                                               116826-20-1
                                                             116826-21-2
     116826-22-3
                  116826-23-4
        (charge-transporting layer contg., for electrophotog.
        photoreceptor)
     116826-22-3
ΙΤ
        (charge-transporting layer contg., for electrophotog.
        photoreceptor)
     116826-22-3 ZCAPLUS
RN
     Benzaldehyde, 4,4'-[1,5-pentanediylbis(phenylimino)]bis[2-methyl-,
CN
     bis[(3-methyl-2(3H)-benzothiazolylidene)hydrazone] (9CI) (CA INDEX
```

NAME)

Me N-N-CH
$$\sim$$
 N- (CH₂) 5-N \sim CH \sim N-N-Me

PAGE 1-B

L9 ANSWER 5 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN

AN 1986:139257 ZCAPLUS

DN 104:139257

ED Entered STN: 19 Apr 1986

TI Electrophotographic photoreceptors

IN Watarai, Osamu; Horie, Seiji

PA Fuji Photo Film Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 21 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM G03G005-06

ICA C09B026-02; H01L031-08

CC 74-3 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

FAN.CNT 1

11111	PATENT NO.		DATE	APPLICATION NO.	DATE	
PI	JP 60186847	A2	19850924	JP 1984-42370	198403	
	JP 04005382	В4	19920131		06	

```
Α
                                19860610
                                            US 1985-708461
     US 4594304
                                                                   198503
                                                                   05
                          Α
                                19840306
PRAI JP 1984-42370
CLASS
 PATENT NO.
                 CLASS PATENT FAMILY CLASSIFICATION CODES
 JP 60186847
                 ICM
                        G03G005-06
                        C09B026-02; H01L031-08
                 ICA
     For diagram(s), see printed CA Issue.
GΙ
     Electrophotog. photoreceptors contain .gtoreq.1 hydrazone compd.
AB
     selected from I, II, and III [R, Rl = Cl-12 alkyl, C7-20 aralkyl,
     condensed aryl (2-4 rings); RR1 in combination may complete a
     heterocycle; R2 = H, C1-12 alkyl, C7-20 aralkyl, aryl; R3, R4, R7,
     R8, R9 = C1-12 alkyl, C7-20 aralkyl, aryl; R3R4 in combination may
     complete a heterocycle; R5, R6 = H, C1-12 alkyl, C7-20 aralkyl,
     aryl, halo, alkoxy, aryloxy; A = benzene on naphthalene ring; Z =
     IV; m = 0, 1; n, p = 0-6; R10, R11 = same as R5 and R6; R10R11 may
     combine to form condensed ring; Z1 = 0, S, Se, imino, methylene].
     The hydrazone compds. are esp. useful as electrophotog. charge
     carrier-transporting agents.
     electrophotog charge carrier transporting agent; hydrazone charge
ST
     carrier transporting agent
     Photography, electro-, developers
ΙΤ
        (composite, charge carrier-transporting hydrazone compds. for)
     101158-49-0 101158-50-3 101158-51-4 101158-52-5
ΙT
     101183-43-1
        (electrophotog. charge carrier-transporting agent)
                    101158-35-4P
                                   101158-36-5P
                                                  101158-37-6P
ΙT
     101158-34-3P
     101158-38-7P
                    101158-39-8P
                                   101158-40-1P
                                                  101158-41-2P
                                                  101158-45-6P
                    101158-43-4P
                                   101158-44-5P
     101158-42-3P
     101158-46-7P 101158-47-8P
                                   101158-48-9P
        (prepn. of, as electrophotog. charge carrier-transporting agent)
ΙT
     29666-92-0
        (reaction of, with bis(methylformylanilino)hexane)
     101158-53-6
IT
        (reaction of, with diphenylhydrazine hydrochloride)
ΙT
     101158-51-4
        (electrophotog. charge carrier-transporting agent)
     101158-51-4 ZCAPLUS
RN
     Benzaldehyde, 4,4'-[1,5-pentanediylbis(ethylimino)]bis[2-methyl-,
CN
     bis[(1,3-dihydro-1,3,3-trimethyl-2H-indol-2-ylidene)hydrazone] (9CI)
       (CA INDEX NAME)
```

Me Me Me
$$N-N-CH$$
 $N-CH$ $N-N-CH$ $N-N-CH$ $N-N-CH$ $N-N-CH$ $N-N-CH$ $N-N-CH$ Me Me

PAGE 1-B

L9 ANSWER 6 OF 6 ZCAPLUS COPYRIGHT 2005 ACS on STN

AN 1973:491705 ZCAPLUS

DN 79:91705

ED Entered STN: 12 May 1984

TI Nonconjugated bis(arylidene)isophthalazines

AU Shubina, L. V.; Gotsko, N. V.

CS USSR

SO Vestn. Beloruss. Univ. (1972), 2(3), 40-3 From: Ref. Zh., Khim. 1973, Abstr. No. 5Zh211

DT Journal

LA Russian

CC 25-5 (Noncondensed Aromatic Compounds)

GI For diagram(s), see printed CA Issue.

AB Reaction of 1,3-C6H4(CHO)2 with N2H4.H2O in alc. 2 hr at apprx.20.degree. gave 70% 1,3-C6H4(CH:NNH2)2, which with the appropriate aldehyde gave the title azines (I). Among the I prepd. were the following (Ar, % yield given): Ph, 60; 2-HOC6H4, 83; 2-MeOC6H4, 85; 2,5-HO(O2N)C6H3, 77; 2-naphthyl, 69. HCl salts were prepd. and characterized.

ST isophthalazine bisarylidene; arylideneisophthalazine; azine isophthalaldehyde arom aldehyde

IT Azines

(of arom. aldehydes and isophthalaldehyde)

42546-09-8P 42546-10-1P **42546-11-2P** TΤ 36604-00-9P 42546-14-5P 42546-15-6P 42546-16-7P 42546-12-3P 42546-13-4P (prepn. of) ΙT 42546-17-8 (reaction of, with arom. aldehydes) 97-51-8 100-10-7 104-88-1 123-11-5 135-02-4 ΙT 66-99-9 574-96-9 (reaction of, with isophthalaldehyde dihydrazone) 100-52-7, reactions IT . 90-02-8, reactions (with isophthalaldehyde dihydrazone) ΙT 42546-11-2P (prepn. of) 42546-11-2 ZCAPLUS RN 1,3-Benzenedicarboxaldehyde, bis[[(2-hydroxy-5-CN nitrophenyl)methylene]hydrazone] (9CI) (CA INDEX NAME)

$$\begin{array}{c} OH \\ \hline \\ NO_2 \end{array} \qquad CH \begin{array}{c} OH \\ \hline \\ NNO_2 \end{array} \qquad CH \begin{array}{c} OH \\ \hline \\ NNO_2 \end{array}$$

Ref #	Hits	Search Query	DBs	Default Operator	Piurals	Time Stamp
Li	4338	azine or ketazine	USPAT	OR	OFF	2005/03/04 16:47
L2	1669	azine or ketazine	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:47
13	1495	azine or ketazine and amine	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:47
L4	1492	azine or ketazine and amine and charge	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:47
L5	1492	azine or ketazine and amine and charge adj transport	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:47
L6	1669	azine or ketazine	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 17:05
L7	47	l6 and charge adj transport	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:56
L8	102	I6 and electrophoto\$	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:57
L9	0	18 not 16	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:57
L10	67	18 not 17	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 16:57
L11	1567	16 not 18	EPO; JPO; DERWENT; IBM_TDB	OR	OFF	2005/03/04 17:06